

Philadelphia College of Osteopathic Medicine DigitalCommons@PCOM

Koch Collection, Papers of Richard S Koch, DO

Special Collections

2013

Proving Spinal Visceral Relations in General Practice

Richard S. Koch

Follow this and additional works at: <http://digitalcommons.pcom.edu/koch>



Part of the [Alternative and Complementary Medicine Commons](#), and the [Osteopathic Medicine and Osteopathy Commons](#)

Recommended Citation

Koch, Richard S., "Proving Spinal Visceral Relations in General Practice"(Original Date Unknown, Online Publication Date 2013).
<http://digitalcommons.pcom.edu/koch/38>

This Abdominal and Other Viscera Research is brought to you for free and open access by the Special Collections at DigitalCommons@PCOM. It has been accepted for inclusion in Koch Collection, Papers of Richard S Koch, DO by an authorized administrator of DigitalCommons@PCOM. For more information, please contact library@pcom.edu.

*Summary for
Press Release*

Proving Spinal Visceral Relations
in General Practice

A 15 year medical research project has recently been concluded involving carefully & taken xray studies of 500 patients each attempting to stand straight.

Whereas, on first impression many patients seem to present excellent posture ~~the~~ the vast majority, (96%) actually could not create vertical ~~or~~ straight spines ~~on the back of neck.~~

They instead possessed vertebral segments which deviated quite obviously from ^{the vertical line of} vertical ^{the vertical of} and from each other. These took the forms of curves and lists.

x Insert #1 here

Studies by both fluoroscope, xray, and manual palpation (or ^{"TESTING BY FEEL"} ~~diagnostic procedures~~ of touch testing were likewise made.)

It was established that unlike other unaffected segments these areas suffered either an apparent absence of, or for certain a restricting limitation of the ^{Symmetrical} desirable normal motion usually expected between the vertebrae, ribs and ~~XXXXXXXXXX~~ associated soft tissue *in the average joints.*

Repeated examinations revealed that these asymmetries were daily persistent and in a consistent pattern characteristic of each patient's spine. Furthermore, if not successfully halted or corrected by treatment they became ~~more~~ ^{MORE} progressively pronounced with aging.

Many patients were free of all spinal discomforts for many hours, and days, ^{a month} at a time. However, whenever they did suffer a tiredness, a ache, soreness, or "kink" or "sprain" of the back or neck it repeatedly occurred most commonly in those ^{same} areas of ^{objectively} detectable ~~objective~~ abnormal mechanics and dynamics.

Insert #2
90% of these cases are grossly and distinctly seen to possess lower extremities ^{VARIATIONS} varying ^{LENGTH} 1/8" or more inches in height with some degree of tilting or unleveling of the ^{limb and base} pelvis. The bottom of the spine rests therefore, on a base that is not horizontal. These figures are supported by ^{other} many scientific investigators.

the original
The majority of patients recalled past injuries ~~and~~ ^{or} strains involving the pelvis, back or neck muscles, and joints in the areas of their current distress. ~~These precede the original distress in their cases of chronic recurrent spinal problems by months & yrs.~~
82% of this study group had frequent or occasional symptoms of or findings of some sort of trouble involving some internal organ which received its supply of nerves from these involved segments of the spine. By physical examination, xrays and laboratory evidence the internal condition was determined to be either organic (a "disease") or functional (~~XXXXXXXXXXXXXXXXXXXX~~
~~XXXXXXXXXXXX~~ ^{"ie."} (nerves) a recurrently disturbed function of the organ). ^{possibly an imbalance of the nerves supplying that organ.}
Patients with established organic diseases recalled having had symptoms in their organs for many months or yrs before any consideration was made.
Meticulous questioning of the patient was concerned with minute details that are believed to be generally disregarded in ordinary medical circles. ^{Tidiously Diligent} There was ^{either} search for the presence of any and all symptoms ^{or signs} frequent or only occasional, ~~with tedious diligence.~~